

ICD-9 Crossover to ICD-10

Diagnosis	ICD-9	ICD-10
Anemia, iron	280.9	D50.9
Anemia, unsp	285.9	D64.9
Chronic pain syndrome	338.4	G89.4
Chronic respiratory failure	518.83	J96.10
Coronary artery disease	414	I25.10
Cushing's Syndrome	255	E24.9
Depression	311	F32.9 (F32.0-F32.8)
Diabetes type 1, with hyperglycemia		E10.65
Diabetes type 1, with ketoacidosis without coma		E10.10
Diabetes type 2	250	E11.9
Diabetes type 2, with chronic kidney disease		E11.22
Diabetes type 2, with foot ulcer		E11.62
Diabetes type 2, with hyperglycemia		E11.65
Diabetes type 2, with nephropathy	250.40-250.43	E11.21
Diabetes type 2, with polyneuropathy		E11.42
Diabetes type 2, with unspecified retinopathy		E11.31
Diabetes type 2, without complications		E11.9
		K44.0 with obstruction K44.9
Diaphragmatic hernia	553.3	w.out obs
Dyspepsia	536.8	R10.13+
Elevated Blood pressure	796.2	R03.0
Fatigue	780.79	R53.8
Gastric dumping syndrome	564.2	K91.1
		K21.9 w/out esophagitis
Gastroesophageal reflux	530.81	521.0 with esophagitis
Hyperlipidemia	272.4	E78.5
Hypertensive chronic kidney dz with stage 1-4 CKD		E12.9
Hypertensive heart disease with heart failure		E11.0
Hypertention, essential	401.9	I10
Hypothalamic lesion	253.9	E23.7
Hypothyroidism	244.9	E03.9 **
Hypothyroidism due to surgery		E89.0
Hypothyroidism without goiter	244.0-244.9	E03.1
Insomnia due to sleep apnea	327.01	G47.01 + G47.30
		K58.0 with diarrhea K58.9
Irritable Bowel Syndrome	564.1	w/out diarrhea
Ischemic heart disease, chronic	414.9	I25.9
Morbid obesity	278.01	E66.01 *
Obesity	278	E66.9
Osteoarthritis, primary, hip bilateral	715.95	M16.0
Osteoarthritis, primary, knee, RIGHT	715.96	M17.11
Overweight	278.02	E66.3

Pain, back	724.5	M54.5 Low M54.6 Thoracic M25.551 (R) M25.552 (L)
Pain, hip	719.45	M25.559 unsp M25.561 (R) M25.562 (L)
Pain, knee	719.46	M25.569 unsp
Postsurgical malabsorption	579.3	K91.2
Pulmonary heart disease	416.9	I27.9
Reactive hypoglycemia	251.2	E16.2
Sleep apnea	780.37	G47.30
Sleep apnea, obstructive		G47.33